

# ***Leucothoe vaderotti*, a new Atlantic *Leucothoe* (Crustacea, Amphipoda) belonging to the “*spinicarpa*-clade” (Crustacea, Amphipoda)**

Traudl Krapp-Schickel<sup>1</sup>

<sup>1</sup> *Forschungsmuseum A. Koenig, Adenauerallee 160, D-53113 Bonn, Germany*

Corresponding author: Traudl Krapp-Schickel ([traudl.krapp@uni-bonn.de](mailto:traudl.krapp@uni-bonn.de))

---

Academic editor: A.-N. Lörz | Received 25 July 2017 | Accepted 7 October 2017 | Published 23 January 2018

---

<http://zoobank.org/40BA7AA9-3CDC-4522-B6C9-E5C9406BA92B>

---

**Citation:** Krapp-Schickel T (2018) *Leucothoe vaderotti*, a new Atlantic *Leucothoe* (Crustacea, Amphipoda) belonging to the “*spinicarpa*-clade” (Crustacea, Amphipoda). In: Brix S, Lörz A-N, Stransky B, Svavarsson J (Eds) Amphipoda from the IceAGE-project (Icelandic marine Animals: Genetics and Ecology). ZooKeys 731: 135–144. <https://doi.org/10.3897/zookeys.731.19813>

---

## **Abstract**

Within the international IceAGE project (Icelandic marine Animals: Genetics and Ecology) some leucothoid amphipods (Crustacea) were collected, among them a rather small new species, belonging to the “*Leucothoe spinicarpa*-clade.”

## **Keywords**

Amphipoda, Atlantic Ocean, IceAGE, Leucothoidae, new species, taxonomy

## **Introduction**

Within the IceAGE-collections (a follow-up of BIOICE, see for example Berge and Vader, 1997) some rather small specimens of the genus *Leucothoe* were collected, similar to the ones described in Krapp-Schickel and Vader (2012) from the Norwegian Sea, called “*Leucothoe* aff. *spinicarpa*”. This genus is extremely easy to recognize as such, but its many species are differentiated from each other only by subtle characters, which has traditionally led to a wholesale lumping of many species under just a few names, especially *Leucothoe spinicarpa* (Abildgaard, 1789). In the last years, it has become clear that the genus, with many of its species living commensally, is a very speciose one and there is no doubt a considerable number of as-yet undescribed species to be discovered (cf. Thomas and Klebba 2007).



## Material and methods

Samples were taken during IceAGE1 and IceAGE2 (with research vessels ‘Meteor’ and ‘Poseidon’; see Brix et al. 2014); they were fixed in cold 96% un-denaturated ethanol, sorted on ice, and stored at 0–4 °C after sorting. They were identified, some mounted on slides with Faure’s fluid, and drawn using a Leitz Laborlux microscope. “Inking” was done with a Wacom tablet, following Coleman, 2003.

Acronyms used in the morphological descriptions are as follows:

<b>A 1, 2</b>	antenna 1, 2	<b>Ped</b>	peduncle
<b>Md</b>	mandible	<b>flag</b>	flagellum
<b>acc.</b>	accessory	<b>Pl</b>	pleopod
<b>Mx 1, 2</b>	maxilla 1, 2	<b>Gn 1, 2</b>	gnathopod 1, 2
<b>ad.</b>	adult	<b>T</b>	telson
<b>Mxp</b>	maxilliped	<b>Hd</b>	head
<b>art</b>	article	<b>U 1–3</b>	uropod 1–3
<b>OP</b>	outer plate	<b>IP</b>	inner plate
<b>Cx</b>	coxal plate	<b>UL</b>	upper lip
<b>P 3–7</b>	peraeopod 3–7	<b>LL</b>	lower lip
<b>Ep</b>	epimeral plate		

## Results

### Leucothoidae Dana, 1852

#### Genus *Leucothoe* Leach, 1814

#### *Leucothoe vaderotti* sp. n.

<http://zoobank.org/BB292914-F4F1-451F-829A-4010FBCF5C31>

Figs 1, 2


*Leucothoe* aff. *spinicarpa* Krapp-Schickel & Vader, 2012, 386–388, fig. 3, 4

**Material.** *Holotype* DZMB–HH 56285 (ZMH K–46787): 1 ad. 5.5 mm, 61°59.83'–61°59.26' N, 000°30.40'–000°32.32'E, Poseidon IceAGE 2, POS 456, 24.7.2013, Norwegian Channel, North Sea, St. 867, 302.5–290 m depth, EBS Supra, 300 µm. In alcohol.

**Additional material.** All in alcohol; one slide DZMB-HH52415.

DZMB–HH 52177 (ZMH K–46788): juv. 2 mm; 61°53.79'N–61°53.53'N, 010°13.77'W–010°12.65'W, Poseidon IceAGE 2, POS 456, 29.7.2013, FI Ridge, St. 878–1, 781.4–775.8 m depth, EBS Supra, 300 µm.

DZMB–HH 52021 (ZMH K–46789): 1 es. 4 mm; 60°24.33'N–60°23.70'N, 006°36.91'–006°38.60'W, Poseidon IceAGE 2, POS 456, 29.7.2013, Farøer Channel South-East, St. 876–5, 554.3–674.8 m depth, shell fragments, EBS, 500 µm.



**Figure 1.** *Leucothoe vaderotti* sp. n. **A 1, A 2** antennae **Mx 1, Mx 2** maxillae **Gn 1, Gn 2** gnathopods **Gn 1', Gn 2'** gnathopods enlarged.



DZMB–HH 52593 (ZMHK–46790): 8 es. 3–4 mm; 61°53.79'N–61°53.53'N, 010°13.77'W–010°12.65'W, Poseidon IceAGE 2, POS 456, 29.7.2013, FI Ridge, St. 878–1, 781.4–775.8 m depth.

DZMB–HH 52617 (ZMH K–46791): 1 juv. 2.5 mm; 63°42.53'N–63°42.78'N, 026°23.05'–026°22.53'W, Meteor IceAGE, ME 85–3, 9.9. 2011, South Iceland, Irminger Basin, slope, St. 1086–1, 698.1–678.5 m depth, 730 trawling distance, EBS Supra, 300 µm.

DZMB–HH 52667 (ZMH K–46792): 1 juv. incompl., 2 mm: 61°38.50'N–61°39.24'N, 031°21.37'–031°20.95'W, Meteor IceAGE, ME 85–3, 7.9.2011, South Iceland, Irminger Basin, Deep Sea, St. 1057–1, 2504.7–2531.8 m depth, 1983 trawling distance, EBS Supra, 300 µm.

DZMB–HH 52694 (ZMH K–46793): 3 juv. 2–3 mm; 60°24.33'N–60°23.70'N, 006°36.91'–006°38.60'W, Poseidon IceAGE 2, POS 456, 29.7.2013, Farøer Channel South-East, St. 876–5, 554.3–674.8 m depth, shell fragments, EBS, 500 µm.

DZMB–HH 52415 (ZMH K–46794): 2 es. 2–3 mm; 63°42.10'N–63°42.37'N, 026°23.64'–026°23.46'W, Meteor IceAGE, ME 85–3, 9.9.2011, South Iceland, Irminger Basin, slope, St. 1082–1, 724.4–704.9 m depth, 782 trawling distance, EBS Supra, 300 µm. Slide in Faure's medium, used for illustration.

**Diagnosis.** Eyes oval, dark in ethanol. Mandibular palp long and narrow, art 3 more than half the length of art 2. Cx 1 inferior margin smooth, nearly as long as wide. Gn 1 carpus distal part approximately six times longer than wide, dactylus reaching approximately 1/3 of propodus length. P 3, P 4 with narrow basis, P 5 – P 7 basis oval and broadened, with regularly rounded and finely serrated hind margin. Ep 2 posterodistally with upturned tip, Ep 3 distoposterior corner with blunt, rounded angle.

**Description.** Length 4–6 mm.

*Head.* Anterior margin rounded, anterodistal margin rectangular with rounded corner. Mid-cephalic keel with acute projection. Rostrum small.

Eyes oval. Antennae short, nearly 1/3 of body length, A 1 peduncle art 1 inferodistally with acute tooth, flagellum up to 11 arts, accessory flagellum not seen. A 2 subequal in length to A 1, peduncle art 4 > art 5, flagellum around 6 arts.

*Mouthparts.* Mandible lacking molars, palp 3-articulate, with long lateral and single distal setae. Art 3 with distal seta, incisors strongly dentate. Other mouthparts like in *L. spinicarpa*.

*Peraeon.* Cx 1–Cx 4 relative width 1: 1.2:0.8:1.


Coxa 1 smooth, length and width subequal; anterodistal margin produced, distal margin regularly rounded, facial setae absent.

Gn 1 basis not inflated, carpus distal part narrow, length to width ratio approximately 6:1; propodus straight, palm with fine short spines; dactylus curved, reaching nearly 1/3 of propodus length.

Cx 2 nearly as long as wide, subquadrangular, much wider than Cx 3, smooth; facial setae absent.

Gn 2 basis slightly inflated, on anterior margin some setae of different length; carpus reaching approx. half propodus, distally truncate, setose; propodus distally with





**Figure 2.** *Leucothoe vaderotti* sp. n. **Mxp** maxilliped **P 3, 4, 5, 7** peraeopods **Ep 2, 3** epimeral plates **Us** urosome with epimeral plates, uropods and telson **Us 1, 2** other urosomites **U1, 2, 3** uropods **T** telson.



short, sharp prolongation, palm convex, regularly rounded, with weak mediofacial setal row, with a few submarginal setae; dactylus curved, smooth, reaching more than 2/3 of propodus length.

Cx 3 length greater than its width, smooth, subrectangular with straight margins and rounded corners.

Cx 4 wider than Cx 3, posterior margin shorter than anterior one, somewhat excavate.

P 3, 4 basis narrow, approx. the width of merus; dactylus reaching or surpassing half the length of propodus.

P 5 – 7 similar, basis oval, both margins with fine serration.

*Pleon.* Ep 1 posteroventral corner rounded. Ep 2 posterodistal corner acutely produced, Ep 3 posteroventral corner bluntly rounded.

U 1 – U 3 similar, length regularly diminishing and U 2 not considerably shorter (like in *L. spinicarpa*).

**Etymology.** Wim Vader completed eight decades in February 2017. He was born Dutch but having lived for much more than half of his life in Norway, he will easily guess that I used the Norwegian word *åtti* = eighty for dedicating this Atlantic species to him. Fifty years ago we began our long-lasting and harmonious collaboration, a “golden jubilee” - thank you, Wim!

**Geographical distribution.** South Iceland- Farøer Channel and Ridge; depth 554–2531 m.

**Remarks.** Together with five specimens collected 1983 between Greenland and Iceland (see Krapp-Schickel and Vader 2012) the present material of the proposed new species consists of 23 animals belonging undoubtedly to the genus *Leucothoe*. All specimens are between 2 and 5 mm, rarely up to 6 mm long. There are four specimens larger than 10 mm with all characters fitting *Leucothoe spinicarpa*, sampled in similar depths as those of the animals 2–5 mm in size. The larger specimens show some differences which are not very conspicuous. It is most probable that they belong to two different species, and it seems also quite reasonable to presume that within the period of early June to end of September these 23 specimens are not all juveniles; however, no ovigerous females were found.

Differences of the small animals compared with *Leucothoe spinicarpa* are:

- Ep 3 totally blunt (see here fig. 2 and Krapp-Schickel and Vader 2012: 390 fig. 4 Ep3) vs. not rounded, but with small but clear posterodistal corner in *L. spinicarpa* (Crowe 2006: 61 fig. 1a, 63 fig. 3e; Sars 1885 pl. 101)
- U 2 in adults reaching length of U1 (see fig. 2) vs. clearly much shorter than U1 and U3 in *L. spinicarpa* (Crowe 2006 fig. 1 and fig. 4)
- Gn 1, 2 basis anterior margin with few irregular longer or shorter setae (see here fig. 1, Krapp-Schickel and Vader 2012: 389 fig. 3) vs. dense setation in *L. spinicarpa* (Crowe 2006 fig. 61 fig. 1 b–e; Sars 1895 pl. 100)
- Gn 1 dactylus less than half the length of propodus (see above fig. 1; Krapp-Schickel and Vader 2012: 389 fig. 3) vs. more than half the length of propodus in *L. spinicarpa* (Crowe 2006 fig. 61 fig 1c; Sars 1885 pl. 100)



- sizes of 23 specimens collected between 1 June–30 September are all between 2–4 mm, with only a few reaching 6 mm; vs. size range between 10–19 mm of *L. spinicarpa* (Crowe 2006, Sars 1895, Krapp-Schickel and Menioui 2005).

These differences are significant enough to distinguish these small specimens as a new species.

**Other material examined.** Two other *Leucothoe* species were sampled, *L. spinicarpa* and *L. lilljeborgi*, the latter clearly in lower depths than the new species.

DZMB–HH 56264 (ZMH K–46795): 1 spec. 2.5 mm *Leucothoe lilljeborgi* 61°25.63'–61°25.05'N, 001°21.07'–001°21.66'E, Poseidon IceAGE 2, POS 456, 24.07.2013, Norwegian Channel, North Sea, St. 866, 169.1–168.8 m depth, EBS Epi, 500 µm.

DZMB–HH 56326 (ZMH K–46796): 2 spec. 2–2.5 mm *L. lilljeborgi* 61°25.63'–61°25.05'N, 001°21.07'–001°21.66'E, Poseidon IceAGE 2, POS 456, 24.7.2013, Norwegian Channel, North Sea, St. 866, 169.1–168.8 m depth, EBS Epi, 300 µm decant.

DZMB–HH 56428 (ZMH K–46797): 1 spec. 2.5 mm *L. lilljeborgi*: 61°25.63'–61°25.05'N, 001°21.07'–001°21.66'E, Poseidon IceAGE 2, POS 456, 24.7.2013, Norwegian Channel, North Sea, St. 866, 169.1–168.8 m depth, EBS Epi, 300 µm.

DZMB–HH 56500 (ZMH K–46798): 1 *L. spinicarpa* 13 mm, 2 males *L. lilljeborgi* 5.5 and 4 mm 61°59.83'–61°59.26'N, 000°30.40'–000°32.32'E, Poseidon IceAGE 2, POS 456, 24.7.2013, Norwegian Channel, North Sea, St. 867, 302.5–290 m depth, EBS Epi, 500 µm.

DZMB–HH 52362 (ZMH K–46799): 1 *L. spinicarpa* 11 mm; 63°42.53'N–63°42.78'N, 026°23.05'–026°22.53'W, Meteor IceAGE, ME 85-3, 9.9.2011, South Iceland, Irminger Basin, slope, St. 1086-1, 698.1–678.5 m depth, 730 trawling distance, EBS Supra, 300 µm.

DZMB–HH 52627 (ZMH K–46800): 1 *L. spinicarpa* 8 mm 62°33.10'N–62°33.22'N, 020°23.71'–020°22.87'W, Meteor IceAGE, ME 85-3, 2.9.2011, South Iceland, Iceland Basin, slope, St. 1010-1, 1384.8–1389 m depth, 1183 trawling distance, EBS Supra, 300 µm.

DZMB–HH 32864 (ZMH K–46801): 1 juv. *Leucothoe* sp. imperf. 2 mm, together with *Leucothoe spinicarpa* 12 mm; 61°53.79'N–61°53.53'N, 010°13.77'W–010°12.65'W, Poseidon IceAGE 2, POS 456, 29.7.2013, FI Ridge, St. 878-1, 781.4–775.8 m depth, EBS Supra, 300 µm.

## Discussion

After Krapp-Schickel and Vader (2012) many *Leucothoe* species are known from the Atlantic Ocean, but mainly from warmer regions. Not much is known about their biology, but it is known that they prefer to live near, in, or with other species such as sponges, and this may also be the reason that they are often well hidden and thus overlooked when generalised ship sampling occurs.



table 1. srekj

Locality	Species	Author
West Africa	<i>Leucothoe minima</i>	Schellenberg, 1925
	<i>L. occidentalis</i>	Reid, 1951
	<i>L. brunonis</i>	Krapp-Schickel & Menioui, 2005
	<i>L. campi</i>	Mateus & Mateus, 1986
	<i>L. spinulosa</i>	Chevreurx, 1920
South Africa	<i>L. miersi</i>	Stebbing, 1888
	<i>L. ctenocheir</i>	K.H. Barnard, 1925
	<i>L. dolichoceras</i>	K.H. Barnard, 1916
Caribbean, Gulf of Mexico	<i>L. wuriti</i>	Thomas & Klebba, 2007
	<i>L. flammosa</i>	Thomas & Klebba, 2007
	<i>L. hortapugai</i>	Winfried et al. 2009
	<i>L. barana</i>	Thomas & Klebba, 2007
	<i>L. ortizi</i>	Winfried et al. 2009
	<i>L. hendrickxi</i>	Winfried et al. 2009
	<i>L. saron</i>	Thomas & Klebba, 2007
	<i>L. ubouhu</i>	Thomas & Klebba, 2006
	<i>L. garigunae</i>	Thomas & Klebba, 2007
Brazil	<i>L. cheiriserra</i>	Serejo, 1998
	<i>L. libue</i>	Barnard, 1970
	<i>L. basilobata</i>	Serejo, 1998
	<i>L. urospinosa</i>	Serejo, 1998
	<i>L. leptosa</i>	Serejo, 1998
Biscaya and Azores	<i>L. cathalaa</i>	Frutos & Sorbe, 2012
	<i>L. rostrata</i>	Chevreurx, 1908
Barbados, Mid Atlantic ridge	<i>L. ayrtonia</i>	Bellan-Santini, 1997
	<i>L. atosi</i>	Bellan-Santini, 1997
Great Britain, Scotland, northern Atlantic	<i>L. incisa</i>	Robertson, 1892
	<i>L. procera</i>	Bate, 1857
	<i>L. lilljeborgi</i>	Boeck, 1861
	<i>L. richiardii</i>	Lessona, 1865
	<i>L. spinicarpa</i>	Abildgaard, 1789
	<i>L. articulosa</i>	(Montagu, 1804)
	<i>L. uschakovi</i>	Gurjanova, 1951

Acknowledgements

I want to thank the IceAGE community for cheerful cooperation, friendship, and many unforgettable hours spent together.

References

Abildgaard PC (1789) Zoologia Danica seu animalium Daniae et Norvegiae rariorum ac minus notorum. Descriptiones et Historia. Havniae, N. Möller et filius, 71 pp. [pls 81–120]



- Barnard KH (1916) Contributions to the crustacean fauna of South Africa. 5. The Amphipoda. Annals of the South African Museum 15(3): 105–302. [pls 26–28] <https://doi.org/10.5962/bhl.title.10646>
- Barnard KH 1925 Contributions to the crustacean fauna of South Africa No. 8 – Further additions to the list of Amphipoda. Annals of the South African Museum 20: 319–380.
- Bate CS (1857) A synopsis of the British edriophthalmous Crustacea. Part I. Amphipoda. Annals and magazine of natural history (2)19, 110: 135–152. [2 figs]
- Bate CS (1862) Catalogue of the specimens of Amphipodous Crustacea in the collections of the British Museum. Taylor & Francis, London, 399 pp. [2–58 pls]
- Bellan Santini D (1997) Amphipods of the cold seep community on the South Barbados accretionary prism. Crustaceana 70(1): 1–30. <https://doi.org/10.1163/156854097X00311>
- Berge J, Vader W (1997) Stegocephalid (Crustacea, Amphipoda) species collected in the BIOFAR and BIOICE programmes. Sarsia 82(4): 347–370. <http://dx.doi.org/10.1080/00364827.1997.10413662>
- Boeck A (1861) Bemaerkninger angaaende de ved de norske kyster forekommende Amphipoder. Forhandlinger ved de Skandinaviske naturforskeres ottende møde i København 8: 631–677.
- Brix S, Meissner K, Stransky B et al. (2014) The IceAge project – a follow up of BIOICE. Polish Polar Research 35(2): 141–150. <https://doi.org/10.2478/popore-2014-0010>
- Chevreaux E (1908) Diagnoses d'amphipodes nouveaux provenant des campagnes de la Princesse-Alice dans l'Atlantique nord. Bulletin de l'Institut Océanographique de Monaco 122: 1–8. [4 figs]
- Chevreaux E (1920) Sur quelques Amphipodes nouveaux ou peu connus provenant des côtes de Bretagne. Bulletin de la Société zoologique de France 45: 75–87.
- Crowe SE (2006) A redescription of *Leucothoe spinicarpa* (Abildgaard, 1789) based on material from the North Atlantic (Amphipoda, Leucothoidae). Zootaxa 1170: 57–68.
- Frutos I, Sorbe C (2012) *Leucothoe cathalaa* sp. nov. (Crustacea: Amphipoda: Leucothoidae), a new bathyal benthic species from the Le Danois Bank ('El Cachucho' Spanish MPA), southern Bay of Biscay. Journal of the Marine Biological Association of the United Kingdom 93(3): 659–666. doi:10.1017/S0025315412000574
- Gurjanova E (1951) Amphipoda Gammaridea from the seas of the USSR and vicinity. Opre-deliteli Faune SSSR Zool Inst Akad Nauk SSSR 41: 1–1029. [In Russian]
- Krapp-Schickel T, Menioui M (2005) *Leucothoe* species from Moroccan Atlantic coasts with redefinition of some species within the *Leucothoe spinicarpa* clade. Bollettino Museo Civico Storia Naturale Verona Botanica-Zoologia 29: 63–83.
- Krapp-Schickel T, Vader W (2012) Leucothoid and Maerid Amphipods (Crustacea) from deep regions of the North Atlantic. Helgoland Marine Research 67(2): 383–396. <https://doi.org/10.1007/s10152-012-0330-3>
- Lessona M (1865) Sopra due nuove specie di animali invertebrati raccolte nel Golfo di Genova. Atti della Società Italiana di Scienze Naturali 8: 423–428.
- Mateus A, Mateus E (1986) Campagne de la 'Calypso' dans le Golfe de Guinée et aux Îles Principe, Sao Tomé et Annobon (1956). Amphipodes récoltés à bord de la 'Calypso'. Anais Faculdade Ciencias Porto 66: 125–223.



- Montagu G (1804) Description of several marine animals found on the south coast of Devonshire. Transaction Linnean Society London 7: 61–85. <https://doi.org/10.1111/j.1096-3642.1804.tb00282.x>
- Reid DM (1951) Report on the Amphipoda (Gammaridea and Caprellidea) of the coast of tropical West Africa. Atlantide Report 2: 189–291.
- Robertson D (1892) A second contribution towards a catalogue of the Amphipoda and Isopoda of the Firth of Clyde and west of Scotland. Proceedings and Transactions of the Natural History Society of Glasgow 3: 199–223.
- Sars GO (1885) Zoology. Crustacea I. Norke Nordhavs Expedition 1876–1878, zoologi 4, 280 pp. [21 pls]
- Sars GO (1890–95) Amphipoda. An account of the Crustacea of Norway with short descriptions and figures of all the species. I. Alb. Cammermeyer, Christiania & Copenhagen, 711 pp [vol. 7], 11 pp [vol. 9]. [+ 240–V111 pl]
- Schellenberg A (1925) Crustacea : Amphipoda. Beiträge zur Kenntnis der Meeresfauna Westafrikas, W. Michaelsen 3, 4: 113–204. [27 fig.]
- Serejo CS (1998) The genus *Leucothoe* (Crustacea, Amphipoda, Leucothoidae) on the Brazilian coast. Beaufortia 48(6): 105–135.
- Stebbing TRR (1888) Report on the Amphipoda collected by H.M.S. Challenger during the years 1873–1876. Eyre and Spottiswoodie, London, Zoology 29: 1–1737.
- Thomas JD, Klebba KN (2006) Studies of commensal Leucothoid Amphipods: Two new sponge-inhabiting species from South Florida and the Western Caribbean. Journal of Crustacean Biology 26(1): 13–22. <https://doi.org/10.1651/C-2624.1>
- Thomas JD, Klebba KN (2007) New species and host associations of commensal leucothoid amphipods from coral reefs in Florida and Belize. Zootaxa 1494: 1–44. <https://doi.org/10.11646/zootaxa.1400.1.1>
- Thomas JD, Ortiz M (1995) *Leucothoe laurensi*, a new species of leucothoid Amphipod from Cuban waters (Crustacea: Amphipoda: Leucothoidae). Proceedings of the biological Society of Washington 108: 613–616.
- Winfield I, Alvarez F (2009) Two new species of amphipods (Peracarida, Amphipoda, Leucothoidae) from the Veracruz Coral Reef system, SW Gulf of Mexico. Crustaceana 82(1): 11–25. <https://doi.org/10.1163/156854008X36721412>